


1/14



IT9302 inhibits spontaneous IL-8 production by purified cultured monocytes. (▲) Indicates the level of IL-8 when using rh IL-10 (100 ng/ml).

Fig. 1

2/14

IT9302 inhibits IL-1 induced (1 ng/ml) IL-8 production by human peripheral blood mononuclear cells




Fig. 2

3/14

## IT9302 stimulated human monocytes




Fig. 3

4/14

# IL-10 stimulated human monocytes




Fig. 4

5/14

Chemotactic activity of IT9302  
on CD8+ T cells




Fig. 5

6/14

Desensitization of CD8+ T cells by  
 IT9302, resulting in unresponsiveness  
 of CD8+ T cells towards IL-10 (10 ng/ml) induced chemotaxis




Fig. 6

7/14

# Suppression of IL-8 activity by IT9302

Effect of IT9302 on IL-8 (10 ng/ml) mediated chemotaxis.




Fig. 7

8/14

IT9302 inhibits MCAF/MCP-1  
induced monocyte chemotaxis





Fig. 8



9/14

ECL-Western Blotting of CD4+ T cell cytosolic proteins using a goat anti-human IL-4 antibody.




Stimulation for 3 days with:

1. Control
2. mon.anti-IL-8 antibody WS.4 10 µg/ml,
3. rIL-8 100 ng/ml,
4. rIL-10 100 ng/ml,
5. IT9302 10 ng/ml,
6. rIFN gamma 10 ng/ml.

Fig. 9

10/14

ECL-Western Blotting of Human Mixed lymphocyte culture cytosolic proteins using a rabbit anti-human TNF- $\alpha$  antibody.



Stimulation for 24 hours with:

1. Control
2. IT9302 10 ng/ml

Fig. 10

11/14

# Regulation of T cell proliferation by IL-10 and IT9302





Fig. 11

12/14

HLA-DR expression on human monocytes (Flow-cytometry):



Non-stimulated cells      + IFN-Gamma (10 ng/ml)      +IL-10 (10 ng/ml)+IFN(10 ng/ml)      +IT9302 (1 ng/ml)+IFN(10 ng/ml)

% cells expressing HLA-DR:

36.8

58.4

25.2

31.2

% cells in apoptosis:

6.6

4.1

10.3

9.3

Fig. 12

13/14




Fig. 13

Leukocyte count in peripheral blood (rabbits)  
after induction of acute pancreatitis (5% bile add)




Fig. 14